

## Pan-Canadian Pharmaceutical Alliance: Completed Negotiations

As of September 30, 2019

321 joint negotiations have been **completed\*\*** for the following drugs and indications:

Drug Product Brand Name (Generic Name)	Indication/Use <i>Please refer to individual jurisdictions for specific reimbursement criteria</i>
<b>Abilify Maintena (aripiprazole long acting injection)</b>	Used for the maintenance treatment of schizophrenia
<b>Abraxane (nab-paclitaxel)</b>	Used first-line to treat locally advanced unresectable or metastatic adenocarcinoma of the pancreas
<b>Actemra (tocilizumab)</b>	Used to treat giant cell arteritis
<b>Actemra (tocilizumab) SC</b>	Used to treat rheumatoid arthritis
<b>Actikerall (fluorouracil and salicylic acid)</b>	Used to treat hyperkeratotic actinic keratosis
<b>Adcetris (brentuximab)</b>	Used to treat lymphoma
<b>Adcetris (brentuximab vedotin)</b>	Used to treat Hodgkin's lymphoma
<b>Adempas (riociguat)</b>	Used to treat chronic thromboembolic pulmonary hypertension
<b>Advair (salmeterol xinafoate/fluticasone propionate)</b>	Used to treat chronic obstructive pulmonary disease and asthma
<b>Afinitor (everolimus)</b>	Used to treat pancreatic neuroendocrine tumours, and to treat breast cancer
<b>Afinitor (everolimus)</b>	Used to treat neuroendocrine tumours of gastrointestinal or lung origin
<b>Akynzeo (netupitant/palonosetron)</b>	Used for the prevention of chemotherapy induced nausea and vomiting
<b>Alecensaro (alectinib)</b>	Used first line to treat locally advanced or metastatic non-small cell lung cancer
<b>Alecensaro (alectinib)</b>	Used second line to treat locally advanced or metastatic non-small cell lung cancer
<b>Alimta (pemetrexed)</b>	Used to treat lung cancer (new indication)
<b>Anoro Ellipta (umeclidinium bromide/vilanterol trifenate)</b>	Used to treat chronic obstructive pulmonary disease
<b>Aptiom (eslicarbazepine)</b>	Used to treat seizures
<b>Arnuity Ellipta (fluticasone furoate)</b>	Used to treat asthma
<b>Aubagio (teriflunomide)</b>	Used to treat relapsing-remitting multiple sclerosis
<b>Avastin (bevacizumab)</b>	Used to treat persistent, recurrent, or metastatic cervical cancer Used to treat epithelial ovarian cancer
<b>Avastin (bevacizumab)</b>	Used to treat genitourinary platinum-resistant ovarian cancer
<b>Avastin (bevacizumab)</b>	Used to treat metastatic colorectal cancer
<b>Basaglar (insulin glargine)</b>	Used to treat Type 1 and Type 2 diabetes mellitus
<b>Bavencio (avelumab)</b>	Used to treat metastatic Merkel cell carcinoma
<b>Besponsa (inotuzumab ozogamicin)</b>	Used to treat acute lymphoblastic leukemia
<b>Biktarvy (bictegravir/emtricitabine/tenofovir alafenamide)</b>	Used to treat HIV infection
<b>Blincyto (blinatumomab)</b>	Used to treat acute lymphoblastic leukemia
<b>Blincyto (blinatumomab)</b>	Used to treat adult acute lymphoblastic leukemia
<b>Blincyto (blinatumomab)</b>	Used to treat pediatric acute lymphoblastic leukemia
<b>Blincyto (blinatumomab)*</b>	Used to treat Philadelphia chromosome positive (Ph+) B-cell precursor (BCP) acute lymphoblastic leukemia (ALL)
<b>Bosulif (bosutinib)</b>	Used to treat chronic myelogenous leukemia (CML)

<b>Botox (onabotulinumtoxin A)</b>	Used to treat multiple indications
<b>Brenzys (etanercept)</b>	Used to treat rheumatoid arthritis and ankylosing spondylitis
<b>Breo Ellipta (fluticasone furoate/vilanterol)</b>	Used for the treatment of asthma
<b>Breo Ellipta (fluticasone furoate/vilanterol)</b>	Used to treat chronic obstructive pulmonary disease
<b>Brilinta (ticagrelor)<sup>†</sup></b>	Used with ASA to prevent atherothrombotic events in patients with acute coronary syndrome
<b>Brilinta (ticagrelor)</b>	Used for secondary prevention of atherothrombotic events
<b>Brivlera (brivaracetam)</b>	Used to treat partial-onset seizures
<b>Cimzia (certolizumab)</b>	Used to treat ankylosing spondylitis, psoriatic arthritis
<b>Caprelsa (vandetanib)</b>	Used to treat medullary thyroid cancer
<b>Cosentyx (secukinumab)</b>	Used to treat moderate to severe plaque psoriasis
<b>Cosentyx (secukinumab)</b>	Used to treat psoriatic arthritis
<b>Cosentyx (secukinumab)</b>	Used to treat ankylosing spondylitis
<b>Cotellic (cobimetinib)</b>	Used to treat metastatic melanoma
<b>Cyramza (ramucirumab)</b>	Used to treat gastrointestinal metastatic gastric cancer or gastro-esophageal junction adenocarcinoma
<b>Cystadrops (cysteamine)</b>	Used to treat corneal cystine crystal deposits
<b>Daklinza (daclatasvir)</b>	Used to treat chronic Hepatitis C infection
<b>Darzalex (daratumumab)</b>	Used to treat multiple myeloma
<b>Diacomit (stiripentol)</b>	Used in patients with severe myoclonic epilepsy in infancy
<b>Dificid (fidaxomicin)</b>	Used to treat Clostridium difficile
<b>Duaklir Genuair (acclidinium bromide/formoterol fumarate dehydrate)</b>	Used to treat chronic obstructive pulmonary disease (COPD)
<b>Duodopa (levodopa/carbidopa)</b>	Used to treat Parkinson's Disease
<b>Dupixent (dupilumab)*</b>	Used to treat atopic dermatitis
<b>Dysport Therapeutic (abobotulinumtoxin A)</b>	Used to treat cervical dystonia
<b>Dysport Therapeutic (abobotulinumtoxin A)</b>	Used to treat upper limb spasticity
<b>Dysport Therapeutic (abobotulinumtoxin A)</b>	Used for the symptomatic treatment of lower limb spasticity in pediatric patients
<b>Effient (prasugrel)</b>	Used with ASA to prevent atherothrombotic events in patients with acute coronary syndrome
<b>Elelyso (taliglucerase alfa)</b>	Used to treat Gaucher Disease Type 1
<b>Eliquis (apixaban)</b>	Used to prevent strokes in patients with atrial fibrillation, and to prevent deep vein thrombosis
<b>Eliquis (apixaban)</b>	Used to treat venous thromboembolic events and prevent recurrent deep-vein thrombosis and pulmonary embolism
<b>Ella (ulipristal acetate)</b>	Used for the prevention of pregnancy
<b>Enstilar (calcipotriol/betamethasone)</b>	Used to treat psoriasis
<b>Entresto (sacubitril/valsartan)</b>	Used to treat heart failure
<b>Entyvio (vedolizumab)</b>	Used to treat ulcerative colitis
<b>Entyvio (vedolizumab)</b>	Used to treat Crohn's disease
<b>Epclusa (sofosbuvir/velpatasvir)</b>	Used to treat chronic hepatitis C infection
<b>Erelzi (etanercept)</b>	Used to treat ankylosing spondylitis, polyarticular juvenile idiopathic arthritis, and rheumatoid arthritis
<b>Erivedge (vismodegib)</b>	Used to treat metastatic basal cell carcinoma
<b>Erleada (apalutamide)</b>	Used to treat non-metastatic castrate resistant prostate cancer
<b>Esbriet (pirfenidone)</b>	Used to treat idiopathic pulmonary fibrosis
<b>Esbriet (pirfenidone)</b> (second letter of intent)	Used to treat idiopathic pulmonary fibrosis

<b>Eylea (aflibercept)</b>	Used to treat neovascular (wet) age-related macular degeneration (AMD), diabetic macular edema (DME), and/or macular edema secondary to central retinal vein occlusion (CRVO)
<b>Eylea (aflibercept)</b>	Used to treat macular edema secondary to branch retinal vein occlusion
<b>Fabrazyme (agalsidase beta)</b>	Used to treat Fabry disease
<b>Fabrazyme (agalsidase beta)</b> <i>(second letter of intent)</i>	Used to treat Fabry disease
<b>Fasenra (benralizumab)</b>	Used to treat eosinophilic asthma
<b>Ferriprox (deferiprone)</b>	Used to treat Transfusional Iron Overload
<b>Fibrystal (ulipristal acetate)</b>	Used for intermittent treatment of moderate to severe signs and symptoms of uterine fibroids
<b>Fibrystal (ulipristal acetate)</b>	Used to treat uterine fibroids
<b>Firazyr (icatibant)</b>	Used to treat acute attacks of hereditary angioedema
<b>Forxiga (dapagliflozin)</b>	Used to treat Type 2 diabetes
<b>Fulphila (pegfilgastrim)*</b>	Used to treat febrile neutropenia in non-myeloid malignancies
<b>Fycompa (perampanel)</b>	Used to treat epilepsy, primary generalized tonic-clonic seizures
<b>Galafold (migalastat)</b>	Used to treat Fabry Disease
<b>Galexos (simeprevir)</b>	Used to treat chronic hepatitis C Genotype 1 infection
<b>Gazyva (obinutuzumab)</b>	Used to treat chronic lymphocytic leukemia
<b>Gazyva (obinutuzumab)</b>	Used to treat follicular lymphoma
<b>Genvoya (elvitegravir/cobicistat/emtricitabine/tenofovir alafenamide)</b>	Used to treat HIV-1 Infection
<b>Gilenya (fingolimod)<sup>†</sup></b>	Used to treat relapsing-remitting multiple sclerosis
<b>Giotrif (afatinib)</b>	Used to treat advanced non-small cell lung cancer
<b>Giotrif (afatinib)</b> <i>(second letter of intent)</i>	Used to treat advanced non-small cell lung cancer
<b>Glatect (glatiramer acetate)</b>	Used to treat relapsing-remitting multiple sclerosis
<b>Grastofil (filgrastim)</b>	Used for prevention or treatment of neutropenia in various indications
<b>Halaven (eribulin)</b>	Used to treat breast cancer
<b>Harvoni (sofosbuvir/ledipasvir)</b>	Used to treat chronic hepatitis C infection
<b>Harvoni (sofosbuvir/ledipasvir)</b> <i>(second letter of intent)</i>	Used to treat chronic hepatitis C infection
<b>Hemangirol (propranolol)</b>	Used to treat infantile hemangioma
<b>Holkira Pak (ombitasvir/paritaprevir/ritonavir and dasabuvir)</b>	Used to treat chronic hepatitis C infection
<b>Humira (adalimumab)</b>	Used to treat ulcerative colitis
<b>Humira (adalimumab)</b>	Used to treat hidradenitis suppurativa
<b>Ibavyr (ribavirin)</b>	Used to treat chronic hepatitis C infection
<b>Ibrance (palbociclib)</b>	Used to treat advanced breast cancer
<b>Iclusig (ponatinib)</b>	Used to treat Chronic Myeloid Leukemia (CML)/ Acute Lymphoblastic Leukemia
<b>Ilaris (canakinumab)</b>	Used to treat systemic juvenile idiopathic arthritis
<b>Imbruvica (ibrutinib)</b>	Used to treat chronic lymphocytic leukemia or small lymphocytic leukemia
<b>Imbruvica (ibrutinib)</b>	Used to treat relapsed/refractory Mantle Cell Lymphoma
<b>Imbruvica (ibrutinib)</b>	Used to treat chronic lymphocytic leukemia and small lymphocytic lymphoma
<b>Imfinzi (durvalumab)</b>	Used to treat non-small cell lung cancer
<b>Increase Ellipta (umeclidinium)</b>	Used to treat chronic obstructive pulmonary disease (COPD)

<b>Inflectra (infliximab)</b>	Used to treat rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis, plaque psoriasis
<b>Inflectra (infliximab)</b>	Used to treat ankylosing spondylitis, plaque psoriasis, psoriatic arthritis, rheumatoid arthritis, Crohn's disease and Ulcerative Colitis
<b>Inlyta (axitinib)</b>	Used to treat metastatic renal cell carcinoma
<b>Inspirolo Respiat (tiotropium/olodaterol)</b>	Used to treat chronic obstructive pulmonary disease (COPD)
<b>Inspra (eplerenone)</b>	Used to treat New York Heart Association (NYHA) class II systolic heart failure
<b>Invega Trinza (paliperidone palmitate)</b>	Used to treat schizophrenia
<b>Invokana (canagliflozin)</b>	Used to treat Type 2 diabetes
<b>Istodax (romidepsin)</b>	Used to treat relapsed/refractory peripheral T-cell lymphoma
<b>Izba (travoprost)</b>	Used to treat open-angle glaucoma
<b>Jadenu (deferasirox)</b>	Used to treat chronic iron overload
<b>Jakavi (ruxolitinib)</b>	Used to treat intermediate to high risk myelofibrosis
<b>Jakavi (ruxolitinib)</b>	Used to treat polycythemia vera
<b>Janumet (sitagliptin/metformin)</b>	Used to treat Type 2 diabetes mellitus
<b>Januvia (sitagliptin)</b>	Used to treat Type 2 diabetes mellitus
<b>Jardiance (empagliflozin)</b>	Used to treat Type 2 diabetes mellitus
<b>Jardiance (empagliflozin)</b>	Used to treat diabetes mellitus (Type 2) for patients with high cardiovascular risk
<b>Jetrea (ocriplasmin)</b>	Used to treat symptomatic vitreomacular adhesion
<b>Jevtana (cabazitaxel)</b>	Used to treat castration resistant metastatic prostate cancer
<b>Juluca (dolutegravir/rilpivirine)</b>	Used to treat HIV-1 infection
<b>Kadcyla (trastuzumab emtansine)</b>	Used to treat HER-2 positive metastatic breast cancer
<b>Kalydeco (ivacaftor)</b>	Used to treat cystic fibrosis
<b>Kalydeco (ivacaftor)</b>	Used to treat cystic fibrosis in patients with CFTR gating mutations or the R117H gating mutation
<b>Kevzara (sarilumab)</b>	Used to treat rheumatoid arthritis
<b>Keytruda (pembrolizumab)</b>	Used to treat metastatic melanoma
<b>Keytruda (pembrolizumab)</b>	Used to treat non-small cell lung cancer (second line or beyond)
<b>Keytruda (pembrolizumab)</b>	Used to treat non-small cell lung cancer (first line)
<b>Kisqali (ribociclib)</b>	Used to treat advanced or metastatic breast cancer
<b>Kuvan (saproterin)<sup>†</sup></b>	Used to treat phenylketonuria
<b>Kyleena (levonorgestrel)</b>	Used for the prevention of pregnancy
<b>Kyprolis (carfilzomib)</b>	Used to treat multiple myeloma
<b>Kyprolis (carfilzomib)</b>	Used to treat multiple myeloma (relapsed)
<b>Lancora (ivabradine)</b>	Used to treat NYHA class II or III heart failure
<b>Lapelga (pegfilgrastim)</b>	Used to treat febrile neutropenia
<b>Lemtrada (alemtuzumab)</b>	Used to treat relapsing-remitting multiple sclerosis
<b>Lenvima (lenvatinib)</b>	Used to treat thyroid cancer
<b>Lixiana (edoxaban)</b>	Used for prevention of stroke and systemic embolism in patients with non-valvular atrial fibrillation
<b>Lixiana (edoxaban)</b>	Used for the prevention and treatment of venous thromboembolism
<b>Lodalis (colesevelam)</b>	Used to treat hypercholesterolemia
<b>Lonsurf (trifluridine/tipiracil)</b>	Used to treat metastatic colorectal cancer.
<b>Lynparza (olaparib)</b>	Used to treat ovarian cancer
<b>Maviret (glecaprevir/pibrentasvir)</b>	Used to treat hepatitis C

<b>MDK-Nitisinone (nitisinone)</b>	Used to treat hereditary tyrosinemia, type 1
<b>Mekinist (trametinib)</b>	Used to treat metastatic melanoma
<b>Metadol-D (methadone hydrochloride)</b>	Used to treat opioid dependence
<b>Metoject (methotrexate)</b>	Used to treat rheumatoid arthritis
<b>Mictoryl (propiverine)</b>	Used to treat overactive bladder (paediatric)
<b>Mifegymiso (mifepristone/misoprostol)</b>	Used for the medical termination of pregnancy
<b>Movapo (apomorphine hydrochloride)</b>	Used to treat Parkinson's Disease
<b>Mozobil (plerixafor)</b>	Used in combination with <i>granulocyte colony-stimulating factor</i> to support treatment for non-Hodgkin lymphoma and multiple myeloma
<b>Mozobil (plerixafor)</b> <i>(second letter of intent)</i>	Used in combination with <i>granulocyte colony-stimulating factor</i> to support treatment for non-Hodgkin lymphoma and multiple myeloma
<b>Mvasi (bevacizumab)</b>	Used for the treatment of metastatic colorectal cancer
<b>Myrbetriq (mirabegron)</b>	Used to treat overactive bladder
<b>Narcan (naloxone hydrochloride)</b>	Used to treat opioid overdose
<b>Neupro (rotigotine)</b>	Used to treat Parkinson's disease
<b>Nitisinone (nitisinone)</b>	Used to treat hereditary tyrosinemia, type 1
<b>Nucala (mepolizumab)</b>	Used to treat severe eosinophilic asthma
<b>Ocaliva (obeticholic acid)</b>	Used to treat primary biliary cholangitis
<b>Ocrevus (ocrelizumab)</b>	Used to treat primary progressive multiple sclerosis
<b>Ocrevus (ocrelizumab)</b>	Used to treat relapsing-remitting multiple sclerosis
<b>Odefsey (emtricitabine/rilpivirine/tenofovir alafenamide)</b>	Used to treat HIV-1 infection
<b>Ofev (nintedanib)</b>	Used to treat idiopathic pulmonary fibrosis
<b>Onbrez (indacaterol)†</b>	Used as a bronchodilator for chronic obstructive pulmonary disease
<b>Onglyza (saxagliptan)</b>	Used to treat Type 2 diabetes mellitus
<b>Opdivo (nivolumab)*</b>	Used for the adjuvant treatment of melanoma
<b>Opdivo (nivolumab)</b>	Used to treat advanced or metastatic non-small cell lung cancer
<b>Opdivo (nivolumab)</b>	Used to treat metastatic melanoma
<b>Opdivo (nivolumab)</b>	Used to treat advanced or metastatic renal cell carcinoma
<b>Opdivo (nivolumab)</b>	Used to treat squamous cell carcinoma of the head and neck (SCCHN)
<b>Oralair (grass pollen allergen extract)</b>	Used to treat allergic rhinitis with or without conjunctivitis
<b>Orencia (abatacept)</b>	Used to treat rheumatoid arthritis
<b>Orfadin (nitisinone)</b>	Used to treat hereditary tyrosinemia, type 1
<b>Ozempic (semaglutide)</b>	Used to treat Type 2 diabetes mellitus
<b>Perjeta (pertuzumab)</b>	Used to treat breast cancer
<b>Pheburane (sodium phenylbutyrate)</b>	Used to treat urea cycle disorders
<b>Plegridy (PegIFN beta-1a)</b>	Used to treat relapsing-remitting multiple sclerosis (RRMS)
<b>Pomalyst (pomalidomide)</b>	Used to treat multiple myeloma
<b>Pomalyst (pomalidomide)</b> <i>Second letter of intent</i>	Used to treat multiple myeloma
<b>Praluent (alirocumab)</b>	Used for lowering of low density lipoprotein cholesterol (LDL-C) in patients with heterozygous familial hypercholesterolemia (HeFH)
<b>Pradaxa (dabigatran)†</b>	Used to prevent strokes in patients with atrial fibrillation
<b>Prezcobix (darunavir/cobicistat)</b>	Used to treat HIV-1 infection
<b>Probuphine (buprenorphine)</b>	Used to treat of opioid use disorder
<b>Procysbi (cysteamine bitartrate)</b>	Used to treat nephropathic cystinosis

<b>Prolia (denosumab)</b>	Used to treat osteoporosis
<b>Quinsair (levofloxacin)</b>	Used to treat cystic fibrosis with chronic <i>Pseudomonas aeruginosa</i> infections
<b>Ravicti (glycerol phenylbutyrate)</b>	Used to treat urea cycle disorders
<b>Rebif (interferon beta-1a)</b>	Used for clinically isolated syndrome
<b>Remicade (infliximab)</b>	Used to treat multiple indications including rheumatoid arthritis, ulcerative colitis and Crohn's disease
<b>Renflexis (infliximab)</b>	Used to treat rheumatoid arthritis, ankylosing spondylitis, adult Crohn's disease, pediatric Crohn's disease, fistulising Crohn's disease, adult ulcerative colitis, pediatric ulcerative colitis, psoriatic arthritis, plaque psoriasis
<b>Repatha (evolocumab)</b>	Used to treat primary hyperlipidemia and mixed dyslipidemia
<b>Replagal (agalsidase alfa)</b>	Used to treat Fabry Disease
<b>Replagal (agalsidase alfa)</b> <i>(second letter of intent)</i>	Used to treat Fabry Disease
<b>Revestive (teduglutide)</b>	Used to treat short bowel syndrome (SBS)
<b>Revlimid (lenalidomide)</b>	Used for maintenance treatment in multiple myeloma after a stem cell transplant
<b>Revlimid (lenalidomide)</b>	Used to treat newly diagnosed multiple myeloma
<b>Rexulti (brexpiprazole)</b>	Used to treat schizophrenia
<b>Rituxan SC (rituximab)</b>	Used to treat multiple indications
<b>Rituxan SC (rituximab)</b>	Used to treat chronic lymphocytic leukemia
<b>Rydapt (midostaurin)</b>	Used to treat Acute Myeloid Leukemia
<b>Seebri Breezhaler (glycopyrronium bromide)</b>	Used to treat chronic obstructive pulmonary disease
<b>Siliq (brodalumab)</b>	Used to treat moderate to severe plaque psoriasis
<b>Simbrinza (brinzolamide/brimonidine)</b>	Used to treat glaucoma and ocular hypertension
<b>Simponi (golimumab) SC</b>	Used to treat ulcerative colitis
<b>Soliris (eculizumab)<sup>†</sup></b>	Used to treat paroxysmal nocturnal hemoglobinuria
<b>Sovaldi (sofosbuvir)</b>	Used to treat chronic hepatitis C infection
<b>Sovaldi (sofosbuvir)</b> <i>(second letter of intent)</i>	Used to treat chronic hepatitis C infection
<b>Spinraza (nusinersen)</b>	Used to treat spinal muscular atrophy
<b>Spinraza (nusinersen)</b>	Used to treat 5q spinal muscular atrophy
<b>Spiriva Respimat (tiotropium)</b>	Used to treat chronic obstructive pulmonary disease
<b>Stivarga (regorafenib)</b>	Used to treat metastatic and/or unresectable gastrointestinal stromal tumours (GIST)
<b>Stivarga (regorafenib)</b> <i>(second letter of intent)</i>	Used to treat metastatic and/or unresectable gastrointestinal stromal tumours (GIST)
<b>Stivarga (regorafenib)</b>	Used to treat unresectable hepatocellular carcinoma
<b>Strensiq (asfotase alfa)</b>	Used to treat pediatric-onset hypophosphatasia
<b>Stribild (elvitegravir/cobicistat/emtricitabine/tenofovir)</b>	Used to treat HIV-1 infection in antiretroviral treatment-naïve adult patients
<b>Sunvepra (asunaprevir)</b>	Used to treat chronic Hepatitis C
<b>Sutent (sunitinib)</b>	Used to treat pancreatic neuroendocrine tumours
<b>Synjardy (empagliflozin and metformin)</b>	Used to treat Type 2 diabetes mellitus in patients with high cardiovascular risk
<b>Sylvant (siltuximab)</b>	Used to treat multicentric Castleman's disease
<b>Tafinlar (dabrafenib)</b>	Used to treat metastatic melanoma
<b>Tafinlar/Mekinist (dabrafenib/trametinib)</b>	Used to treat melanoma
<b>Tafinlar/Mekinist (dabrafenib/trametinib)</b>	Used to treat metastatic melanoma (in combination)

<b>Tagrisso (osimertinib)</b>	Used to treat locally advanced or metastatic EGFR T790M mutation-positive non-small cell lung cancer (NSCLC) who have progressed on or after EGFR TKI therapy
<b>Taltz (Ixekizumab)</b>	Used to treat moderate to severe plaque psoriasis
<b>Taltz (ixekizumab)</b>	Used to treat psoriatic arthritis
<b>Tecentriq (atezolizumab)</b>	Used to treat non-small cell lung cancer
<b>Tecfidera (dimethyl fumarate)</b>	Used to treat multiple sclerosis
<b>Tivicay (dolutegravir)</b>	Used to treat HIV-1 infection
<b>Tobi Podhaler (tobramycin)</b>	Used for the management of cystic fibrosis patients with chronic pulmonary <i>Pseudomonas aeruginosa</i> infections
<b>Tradjenta (linagliptin) and Jentadueto (linagliptin/metformin)</b>	Used to treat Type 2 diabetes mellitus
<b>Treanda (bendamustine)<sup>†</sup></b>	Used to treat non-Hodgkin lymphoma, mantle cell lymphoma, chronic lymphocytic leukemia
<b>Toujeo (insulin glargine)</b>	Used to treat Type 1 and Type 2 diabetes mellitus
<b>Trelegy Ellipta (fluticasone furoate/umeclidinium/vilanterol)*</b>	Used to treat chronic obstructive pulmonary disease
<b>Tresiba (insulin degludec)</b>	Used to treat Type 1 and Type 2 diabetes mellitus
<b>Triumeq (dolutegravir/abacavir/lamivudine)</b>	Used to treat HIV-1 infection
<b>Tudorza Genuair (aclidinium bromide)</b>	Used in management of chronic obstructive pulmonary disease (COPD)
<b>Tysabri (natalizumab)</b>	Used to treat relapsing-remitting multiple sclerosis
<b>Ultibro (indacaterol/glycopyrronium)</b>	Used to treat chronic obstructive pulmonary disease
<b>Unituxin (dinutuximab)</b>	Used in combination with GM-CSF, IL-2 and Retinoic acid (RA) for the treatment of pediatric patients with high-risk neuroblastoma
<b>Uptravi (selexipag)</b>	Used to treat pulmonary arterial hypertension
<b>Valtrex (valacyclovir)</b>	Used to treat specific viral infections
<b>Vectibix (panitumumab)</b>	Used to treat metastatic colorectal cancer
<b>Velphoro (sucroferric oxyhydroxide)*</b>	Used to treat hyperphosphatemia in patients with end stage renal disease.
<b>Venclexta (venetoclax)</b>	Used to treat chronic lymphocytic leukemia
<b>Vimizim (elosulfase alfa)</b>	Used to treat mucopolysaccharidosis IVA (Morquio A syndrome)
<b>Vosevi (sofosbuvir velpatasvir voxilaprevir)</b>	Used to treat chronic hepatitis C infection
<b>Votrient (pazopanib)</b>	Used to treat metastatic renal cell carcinoma
<b>Votrient (pazopanib) (second letter of intent)</b>	Used to treat metastatic renal cell carcinoma
<b>VPRIV (velaglucerase)</b>	Used to treat Gaucher Disease Type 1
<b>Xalkori (crizotinib)</b>	Used to treat advanced non-small cell lung cancer (second-line)
<b>Xalkori (crizotinib)</b>	Used to treat advanced non-small cell lung cancer (first-line)
<b>Xarelto (rivaroxaban)</b>	New indication for treatment of venous thromboembolic events and prevention of recurrent deep vein thrombosis and pulmonary embolism.
<b>Xarelto (rivaroxaban)<sup>†</sup></b>	Used to prevent strokes in patients with atrial fibrillation, and to prevent deep vein thrombosis
<b>Xeljanz (tofacitinib)</b>	Used to treat rheumatoid arthritis
<b>Xeljanz XR (tofacitinib)</b>	Used to treat rheumatoid arthritis
<b>XigDuo (dapagliflozin/metformin hydrochloride)</b>	Used to treat Type 2 diabetes mellitus
<b>Xolair (omalizumab)</b>	Used to treat chronic idiopathic urticaria
<b>Xtandi (enzalutamide)</b>	Used to treat first-line metastatic castrate resistant prostate cancer (before chemotherapy)
<b>Xtandi (enzulatamide)</b>	Used to treat castrate resistant prostate cancer (after chemotherapy)

<b>Yervoy (ipilimumab)</b>	Used for the first-line treatment of adult patients with advanced (unresectable or metastatic) melanoma
<b>Yervoy (ipilimumab)<sup>†</sup></b>	Used for the second-line treatment of adult patients with unresectable or metastatic melanoma
<b>Yervoy/Opdivo (nivolumab/ipilimumab)</b>	Used to treat metastatic melanoma
<b>Yervoy/Opdivo (nivolumab/ipilimumab)</b>	Used to treat renal cell carcinoma
<b>Zaxine (rifaximin)</b>	Used to treat hepatic encephalopathy
<b>Zepatier (elbasvir grazoprevir)</b>	Used for the treatment of Hepatitis C (genotype 1, 4)
<b>Zirabev (bevacizumab)</b>	Used for the treatment of metastatic colorectal cancer
<b>Zydelig (idelalisib)</b>	Used to treat relapsed chronic lymphocytic leukemia (CLL)
<b>Zykadia (ceritinib)</b>	Used to treat ALK+ non-small cell lung cancer
<b>Zytiga (abiraterone)</b>	Used to treat prostate cancer

#### 45 drug negotiations were closed as agreements were not reached:

<b>Abilify (aripiprazole)</b>	Used for the treatment of schizophrenia and related psychotic disorders
<b>Adempas (riociguat)</b>	Used to treat pulmonary arterial hypertension
<b>Aermony RespiClick (fluticasone propionate)</b>	Used to treat asthma
<b>Arbesda RespiClick (fluticasone propionate/salmeterol xinafoate)</b>	Used to treat asthma
<b>Brilinta (ticagrelor)</b>	Used with ASA to prevent atherothrombotic events with history of myocardial infarction
<b>Butrans (buprenorphine)</b>	Used to treat persistent pain of moderate intensity
<b>Byetta (exenatide)</b>	Used to treat Type 2 diabetes
<b>Campral (acamprosate calcium)</b>	Used for maintenance of alcohol abstinence
<b>Caprelsa (vandetanib)</b>	Used to treat medullary thyroid cancer
<b>Cerdelga (eliglustat)</b>	Used to treat Gaucher Disease Type 1
<b>Cerezyme (imiglucerase)</b>	Used to treat Gaucher Disease Type 1
<b>Cinqair (reslizumab)</b>	Used to treat severe eosinophilic asthma
<b>Copaxone (glatiramer acetate)</b>	Used to treat multiple sclerosis
<b>Daklinza (daclatasvir)</b>	Used to treat chronic Hepatitis C
<b>Descovy (emtricitabine/tenofovir alafenamide)</b>	Used to treat HIV-1 infection
<b>Faslodex (fulvestrant)</b>	Used to treat locally advanced or metastatic breast cancer
<b>Fiasp (insulin aspart)</b>	Used to treat Type 1 diabetes mellitus
<b>Fibristal (ulipristal acetate)</b>	Used to treat uterine fibroids (2nd cycle pre-surgery)
<b>Invokamet (canagliflozin/metformin hydrochloride)</b>	Used to treat Type 2 diabetes mellitus
<b>Kalydeco (ivacaftor)</b>	Used to treat cystic fibrosis with G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N, S549R, G970R or R117H mutation
<b>Kevzara (sarilumab)</b>	Used to treat rheumatoid arthritis
<b>Lartruvo (olaratumab)</b>	Used to treat advanced soft tissue sarcoma
<b>Methadose (methadone)</b>	Used to treat opioid dependence
<b>Mictoryl (propiverine)</b>	Used to treat overactive bladder
<b>Monoprost (latanoprost)</b>	Used to treat glaucoma and ocular hypertension
<b>Onivyde (irinotecan liposome)</b>	Used to treat metastatic pancreatic cancer



<b>Opsumit (macitentan)</b>	Used to treat pulmonary arterial hypertension
<b>Otezla (apremilast)</b>	Used to treat moderate to severe plaque psoriasis
<b>Otezla (apremilast)</b>	Used to treat psoriatic arthritis
<b>Repatha (evolocumab)</b>	Used for the prevention of cardiovascular events in patients with atherosclerotic cardiovascular disease (ASCVD)
<b>Rosiver (ivermectin)</b>	Used to treat rosacea
<b>Rosiver (ivermectin)</b>	Used to treat rosacea
<b>Simponi (golimumab) IV</b>	Used to treat rheumatoid arthritis
<b>Soliris (eculizumab)</b>	Used to treat aHUS
<b>Stelara (ustekinumab)</b>	Used to treat Crohn's disease
<b>Strensiq (asfotase alfa)</b>	Used to treat hypophosphatasia
<b>Symtuza (darunavir/cobicistat/emtricitabine/tenofovir alafenamide)</b>	Used to treat HIV infection
<b>Technivie (ombitasvir/paritaprevir/ritonavir)</b>	Used for the treatment of Hepatitis C (genotype 4)
<b>Tremfya (guselkumab)</b>	Used to treat moderate to severe plaque psoriasis
<b>Trulicity (dulaglutide)*</b>	Used to treat Type 2 diabetes mellitus
<b>Vemlidy (tenofovir alafenamide)</b>	Used to treat chronic hepatitis B infection
<b>Viacoram (perindopril arginine/amlodipine)</b>	Used to treat essential hypertension
<b>Victoza (liraglutide)</b>	Used to treat Type 2 diabetes
<b>Victoza (liraglutide) (second negotiation)</b>	Used to treat Type 2 diabetes mellitus
<b>Xolair (omalizumab)</b>	Used to treat severe persistent asthma

\*\* A completed pan-Canadian negotiation refers to those for which a Letter of Intent (document which outlines the agreed upon terms and conditions for listing) has been signed between the lead jurisdiction for the negotiation and the manufacturer OR negotiations have been closed without a Letter of Intent.

† The initial nine products negotiated through the pCPA.

\* New negotiations that have been completed since last update on August 31, 2019.